


# Normal and Inverse Diffusive Isotope Fractionation of Deuterated Toluene and Benzene in Aqueous Systems

Massimo Rolle\*<sup>†</sup> and Biao Jin<sup>†</sup>

<sup>†</sup>Department of Environmental Engineering, Technical University of Denmark, Bygningstorvet Building 115, DK-2800 Kgs. Lyngby, Denmark

**S** Supporting Information

**ABSTRACT:** Diffusive isotope fractionation of organic contaminants in aqueous solution is difficult to quantify, and only a few experimental data sets are available for compounds of environmental interest. In this study, we investigate diffusive fractionation of perdeuterated and nondeuterated benzene and toluene. Multitracer experiments were carried out in 1-D gel dissection tubes and in a quasi-2-D flow-through porous medium. The experiments allowed us to simultaneously and directly compare the diffusive and dispersive behavior of benzene and toluene. We observed an unexpected, opposite behavior of the two monoaromatic hydrocarbons. Toluene showed a normal diffusive isotope effect ( $D_{C_7D_8}/D_{C_7H_8} = 0.96$ ) with enrichment of the nondeuterated isotopologue in the direction of the diffusive and transverse dispersive fluxes. Conversely, the measured trends for benzene indicate inverse diffusive fractionation ( $D_{C_6D_6}/D_{C_6H_6} = 1.02$ ), with a remarkably faster diffusion rate of the perdeuterated isotopologue that was enriched in the downgradient portion of the diffusion tubes and at the fringes of the contaminant plumes in the flow-through setup. These outcomes can neither be interpreted as mass-dependent fractionation nor be described as purely hydrodynamic (i.e., mass independent) effects. The results of this study are relevant for the use of labeled/nonlabeled mixtures of organic compounds as conservative and (bio)reactive tracers in environmental applications.



## INTRODUCTION

Aromatic hydrocarbons are widespread contaminants, frequently found in soils and aquatic environments. Among these chemicals BTEX (i.e., benzene, toluene, ethylbenzene, and xylene) compounds are of particular concern due to their relatively high solubility, mobility, and toxicity.<sup>1</sup> Mixtures of labeled and nonlabeled compounds have been often applied as a diagnostic tool to understand and quantify contaminant transport and transformation mechanisms. For instance, in the field of groundwater contamination, deuterium-labeled BTEX compounds have been used in tracer tests, push–pull tests, and in situ microcosms to evaluate contaminant retardation and in situ rates of transformation during groundwater bioremediation and natural attenuation.<sup>2–6</sup> One important aspect that has not (yet) been evaluated in detail is the extent of diffusive isotope fractionation of BTEX compounds in aqueous solution and its implications in environmental systems. The evaluation of such an effect is important since recent investigation of subsurface solute transport has highlighted the key controlling role of aqueous diffusion for groundwater contaminant transport at different scales.<sup>7–10</sup> Despite the increased recognition of the quantitative importance and of the macroscopic impact of small-scale diffusive processes on large-scale transport of organic contaminants, only a few experimental and modeling studies have attempted to quantify diffusive isotope fractionation for organic compounds.<sup>11–16</sup> The lack of data and mechanistic understanding of organic chemicals' diffusive

isotope fractionation becomes apparent when compared with the advances in the related field of inorganic isotope geochemistry, in which numerous studies have been carried out to investigate diffusive isotope effects of major cations, anions, and dissolved gases in both aqueous solutions<sup>17–29</sup> and nonaqueous systems.<sup>30–33</sup>


In this work, we focus on the diffusive transport of perdeuterated and nondeuterated benzene and toluene. The diffusive behavior of these chemicals (particularly of benzene) was investigated in early studies,<sup>34–36</sup> mostly as self-diffusion or as tracer diffusion in organic solvents. However, to the best of our knowledge, no study has reported a comparison of diffusive isotope fractionation of these contaminants in aqueous solution. The purpose of this Letter is to report the unexpected, contrasting fractionation behavior of toluene and benzene that we have observed in a series of experiments performed in different setups under purely diffusive conditions (1-D gel dissection tubes), as well as in flow-through systems (2-D flow-through chamber). A key feature of the experiments was the simultaneous presence of nondeuterated and perdeuterated mixtures of toluene and benzene undergoing diffusion and lateral dispersion. In this way, in each experiment, the four

Received: April 29, 2017

Revised: May 29, 2017

Accepted: May 31, 2017

Published: May 31, 2017



**Figure 1.** Spatial profiles of concentrations and isotopologue ratios for benzene and toluene along the length of the diffusion tubes in the 9-day experiment A (a–c) and in the 15-day experiment C (d–f). Symbols (squares for nondeuterated, triangles for perdeuterated isotopologues, circles for ratios) represent the measured data, whereas the lines are the outcomes of the simulations.

tracers ( $C_6H_6$ ,  $C_6D_6$ ,  $C_7H_8$ , and  $C_7D_8$ ) were all transported under the same conditions, which facilitates a direct comparison of the effects of diffusion and transverse dispersion on the concentration of nondeuterated and perdeuterated benzene and toluene and, more importantly, on their diffusive isotope fractionation.

## MATERIALS AND METHODS

**Chemicals and Analytical Methods.** High-purity organic compounds (99.5%, Sigma-Aldrich, Germany) were used in the experiments. Measurements were carried out with a 7890A gas chromatograph (GC) with a capillary column ( $30\text{ m} \times 250\ \mu\text{m}$ ,  $1.0\ \mu\text{m}$  film thickness; Agilent, USA) coupled to a 5975C triaxis mass selective detector (MSD) (Agilent, USA). Headspace samples were injected for analysis using a COMBIPAL multipurpose autosampler system.

**1-D Diffusion Experiment.** Gel dissection experiments were performed in cylindrical glass tubes (1.1 cm diameter and 20 cm length) using agarose gel prepared with a minimum amount (1% w/w) of phyto agar (Duchefa, Netherlands). Agar solutions containing mixtures of dissolved toluene, perdeuterated toluene, benzene, and perdeuterated benzene (1:1:1:1 volume proportion) were prepared and filled in the first 10 cm of the diffusion tubes. This zone acted as contaminant source during the experiments in which the compounds diffused toward the remaining portion of the tubes that was filled with pure gel medium. The tubes were kept horizontally at a

constant temperature of  $20\ ^\circ\text{C}$ . After 9 and 15 days, the tubes were sampled by cutting the gel into 1 cm slices with a scalpel. The slices were immediately sealed in 10 mL glass vials with screw caps and sent to GC analysis.

**2-D Flow-through Experiment.** Flow-through experiments were performed in a quasi-two-dimensional flow-through chamber (inner dimensions:  $80\text{ cm} \times 18\text{ cm} \times 1\text{ cm}$ ,  $L \times H \times W$ ), equipped with 10 equally spaced (1 cm spacing) ports both at the inlet and at the outlet. The flow-through system was filled with homogeneous quartz sand (Euroquartz, Germany) with grain diameter of 1.0–1.5 mm. The sand was washed in an acidic solution and dried for 12 h in an oven at  $120\ ^\circ\text{C}$  before filling the flow-through chamber. The sand was filled with a wet-packing procedure using ultrapure Milli-Q water (Evoqua-Water, USA) to avoid air entrapment in the porous medium.<sup>37</sup> The inlet and outlet ports were connected to two high-precision multichannel peristaltic pumps (IPC-N24, Ismatec, Switzerland). Sampling was performed with a 10-channel syringe pump (KD Scientific, USA). The system was operated at a seepage velocity of 0.8 m/d. An aqueous solution containing the four isotopologues was continuously injected from the two lowermost ports at the inlet of the flow-through chamber. After establishing a steady-state plume (i.e., exchanging at least two pore volumes), samples were taken at the outlet ports and analyzed for the concentrations of deuterated and nondeuterated compounds.

Table 1. Summary of Results for Multitracer 1-D Diffusion Experiments

| experiment     | time [days] | $D_{C_nH_m}$ [ $\times 10^{-9} \text{ m}^2 \text{ s}^{-1}$ ] | $D_{C_nD_m}$ [ $\times 10^{-9} \text{ m}^2 \text{ s}^{-1}$ ] | $\beta$ | $D_{C_nD_m}/D_{C_nH_m}$ |
|----------------|-------------|--|--|---------|-------------------------|
| <b>benzene</b> |             |  |  |         |                         |
| A              | 9           | $0.958 \pm 0.040$  | $0.976 \pm 0.017$  | -0.251  | 1.019                   |
| B              | 9           | $0.958 \pm 0.008$  | $0.978 \pm 0.007$  | -0.279  | 1.021                   |
| C              | 15          | $0.956 \pm 0.021$  | $0.972 \pm 0.020$  | -0.218  | 1.016                   |
| D              | 15          | $0.957 \pm 0.025$  | $0.974 \pm 0.024$  | -0.238  | 1.018                   |
| <b>toluene</b> |             |  |  |         |                         |
| A              | 9           | $0.800 \pm 0.010$  | $0.768 \pm 0.009$  | 0.490   | 0.960                   |
| B              | 9           | $0.811 \pm 0.015$  | $0.780 \pm 0.012$  | 0.460   | 0.962                   |
| C              | 15          | $0.798 \pm 0.012$  | $0.769 \pm 0.010$  | 0.444   | 0.964                   |
| D              | 15          | $0.810 \pm 0.019$  | $0.780 \pm 0.016$  | 0.453   | 0.963                   |

**Modeling Approach.** The governing equations for contaminant transport in the two experimental setups are the 1-D Fick's second law of diffusion<sup>38</sup> in the gel dissection tubes and the 2-D steady state advection dispersion equation in the 2-D flow-through setup.<sup>39,40</sup> The models used to quantitatively interpret the experimental results are based, respectively, on a numerical and an analytical solution of these governing equations. The key parameters controlling transport of the different toluene and benzene isotopologues are their diffusion coefficients and, in the flow-through setup, their transverse hydrodynamic dispersion coefficients. The governing equations and their boundary conditions are summarized in the [Supporting Information](#) together with concentration maps of the steady-state plumes in the 2-D flow-through setup.

## RESULTS AND DISCUSSION

Two examples of benzene and toluene spatial profiles measured in the 1-D setup after 9 and 15 days of diffusion are illustrated in [Figure 1](#). The plots show typical diffusion curves with small differences between deuterated and nondeuterated compounds that are difficult to appreciate in the concentration plots. The differences become apparent by plotting the ratios  $C_6D_6/C_6H_6$  and  $C_7D_8/C_7H_8$ . These ratios surprisingly show an opposite trend with a decreasing pattern for toluene and a reverse, increasing trend for benzene. To quantify these observations, we used a simple inverse power law model,<sup>21,41</sup> which relates the diffusion coefficients of the deuterated and nondeuterated isotopologues to their molecular masses

$$\frac{D_{C_nD_m}}{D_{C_nH_m}} = \left( \frac{m_{C_nH_m}}{m_{C_nD_m}} \right)^\beta \quad (1)$$


Fitting the experimental data allowed estimating the values of the diffusion coefficients for the nondeuterated and D-labeled isotopologues and, thus, the value of the exponent  $\beta$  expressing the mass dependence of the isotopologues diffusion coefficients. The fitting procedure was carried out using the function *lsqnonlin* implemented in MATLAB. [Table 1](#) summarizes the results obtained for the different tube experiments and reports the observed diffusive isotope fractionation. Both the graphical representation of the experimental data ([Figure 1](#)) and the parameters reported in [Table 1](#) show a normal isotope effect for toluene and an inverse isotope effect for benzene. In the case of toluene, the deuterated isotopologue has a slightly lower diffusion coefficient and results in decreasing isotopologue ratios as the toluene species diffuse toward the pure gel medium. For benzene, instead, the data show an inverse diffusive isotope effect: the molecules of the deuterated isotopologue diffuse at a

slightly faster rate and become enriched in the initially pure gel medium as benzene diffuses from the contamination source. This behavior was consistently observed in all experiments. The  $\beta$  (0.444–0.490) values for toluene are consistent with those of previous experiments,<sup>15</sup> whereas the negative values for benzene clearly indicate an inverse isotope effect. Average values characterizing normal and inverse isotope fractionation observed for toluene and benzene in these diffusion experiments are  $D_{C_7D_8}/D_{C_7H_8} = 0.962 \pm 0.002$  and  $D_{C_6D_6}/D_{C_6H_6} = 1.019 \pm 0.002$ .

[Figure 2](#) shows vertical profiles of concentrations and isotopologue ratios observed for the simultaneous transport of deuterated and nondeuterated benzene and toluene in the flow-through chamber. The concentration trends show a typical transverse dispersion profile for plumes continuously injected in a homogeneous porous medium. The isotopologue ratio measured at the different ports confirms the normal diffusive fractionation for toluene and the inverse fractionation for benzene. In fact, the ratio  $C_7D_8/C_7H_8$  decreases from the core toward the outer fringe of the plume, whereas the ratio  $C_6D_6/C_6H_6$  increases toward the plume fringe. No fitting procedure was used to evaluate the data from the flow-through experiment. Instead, the experimental results were evaluated with pure forward modeling of solute transport based on the average diffusion coefficients of the four compounds determined in the tube diffusion setups. The key parameter in this setup is the transverse hydrodynamic dispersion coefficients, which is compound (and isotopologue) specific and was described with the following parameterization:

$$D_T = D_p + D_{aq} \left( \frac{Pe^2}{Pe + 2 + 4\delta^2} \right)^{0.5} \quad (2)$$

where  $D_p$  [ $L^2/T$ ] denotes the velocity-independent pore diffusion coefficient;  $Pe = vd/D_{aq}$  is the grain Péclet number, in which  $d$  [ $L$ ] is the average grain size diameter;  $v$  [ $L/T$ ] is the seepage velocity;  $\delta$  is the ratio between the length of a pore channel and its hydraulic radius. A value of  $\delta = 5.37$  was determined in previous tracer experiments performed in a range of porous media, grain sizes, and seepage velocities comprising the conditions of the current flow-through setup.<sup>42</sup> This parametrization of transverse dispersion has been tested for solute transport in similar quasi-2-D systems<sup>43–45</sup> and verified in pore-scale studies<sup>46,47</sup> and fully three-dimensional flow-through experiments.<sup>42</sup> The essential feature of [eq 2](#) is that it acknowledges the importance of aqueous diffusion also in the nonlinear, velocity-dependent, mechanical dispersion term. The good agreement of the predictive, purely forward simulations for both measured concentrations and ratios ([Figure 2](#)) shows



**Figure 2.** Observed and simulated vertical profiles of concentrations and isotopologue ratios at the outlet of the flow-through setup (a, b, d, e) and spatial maps of the isotopologue ratio for benzene (c) and toluene (f). The concentration data at the outlet ports (squares for the nondeuterated, triangles for the perdeuterated isotopologues) are average of triplicate measurements of two sampling events of the steady-state plumes, carried out after flushing 2 and 4 pore volumes (i.e., 48 and 96 h, respectively).

the capability of the model to capture the diffusive fractionation effects in the flow-through system and also the accuracy of the experimentally determined values of aqueous diffusion for the deuterated and nondeuterated tracers obtained from the 1-D tube experiments. The latter compare very well with the values of aqueous diffusion coefficients computed with classical empirical correlations for organic compounds (Table S2, Supporting Information); this also indicates that sorption of toluene and benzene was not important in our experiments. The values of the transverse dispersion coefficients, determined according to eq 2, and used in the profiles and isotopologue ratio maps shown in Figure 2 are  $D_{T,C7H8} = 1.563 \times 10^{-9}$ ,  $D_{T,C7D8} = 1.547 \times 10^{-9}$ ,  $D_{T,C6H6} = 1.641 \times 10^{-9}$ , and  $D_{T,C6D6} = 1.649 \times 10^{-9}$  m<sup>2</sup>/s.

Diffusive isotope fractionation in condensed systems, such as aqueous solutions, is challenging to investigate and to conceptualize in a solid theoretical framework. Different factors that may affect diffusive isotope fractionation include solute mass, volume, shape, molecular structure, polarity, temperature, and solute–solvent interactions. Mass and volume dependencies have dominated the discussion about diffusive fractionation in condensed systems and the interpretation of diffusion experiments. Models that have been proposed to capture the effects of diffusive fractionation include the so-called hydrodynamic description (e.g., Stokes–Einstein relation), as well as the extension to the condensed phase of formulations derived from Chapman–Enskog kinetic theory of gas diffusion.<sup>38</sup> The hydrodynamic model does not include mass dependence, and the key parameter is the radius, thus, the molar volume of the solute, whereas the kinetic theory predicts an inverse square root relation with the reduced mass.<sup>41</sup> Most

of the experimental studies for different aqueous solutes have reported a mass dependence that is typically weaker than the inverse square root relation from kinetic theory. A simple power law expression (eq 1), in which the exponent is derived by fitting the experimental data, is the model that is most often used to relate the diffusive isotope effects to the mass of the diffusing solutes in theoretical, computational, and experimental studies.<sup>21,24–26,41</sup> It is worth pointing out that in this simple model the exponent  $\beta$  is an empirical coefficient which also lumps other factors such as steric effects and solute–solvent interactions that are likely to play an important role on diffusive isotope fractionation. Indeed, the data presented in this study show normal and inverse isotope effects that can neither be interpreted on the basis of mass dependencies nor be ascribed as purely hydrodynamic (i.e., mass independent) effects. In fact, on the basis of the sole mass difference, the deuterated benzene should have shown lower diffusion rates than the nonlabeled isotopologue. Instead, an inverse effect was observed both in the tube diffusion experiments and in the flow-through setup. Our experiments also allow excluding a purely hydrodynamic interpretation of the results. Molar volumes of deuterated benzene and toluene have been shown to be smaller than those of the corresponding nonlabeled isotopologues; for instance, Bartell and Roskos<sup>48</sup> report the following values at 20 °C:  $V_{C6D6} = 88.62$  mL/mol,  $V_{C6H6} = 88.86$  mL/mol,  $V_{C7D8} = 105.98$  mL/mol, and  $V_{C7H8} = 106.28$  mL/mol. These differences are commonly attributed to the smaller effective radius of the C–D bonds compared to the C–H bonds.<sup>48,49</sup> However, a pure hydrodynamic interpretation, based on a simple Stokes–Einstein relation and considering the molar volumes reported above, would predict very weak inverse isotope effects ( $D_{CnDm}/$

$D_{C_nH_m} = 1.0009$ ) for both compounds. This was not observed in this study, in which stronger inverse and normal diffusive isotope fractionation effects were observed for benzene and toluene, respectively. We hypothesize that the geometry of the hydration shells surrounding the aromatic molecules and the solvation dynamics play a major role for diffusive isotope fractionation of deuterated and nondeuterated benzene and toluene in aqueous solutions. For instance, a recent computational study of benzene hydration<sup>50</sup> reports that the hydration shell of a benzene molecule has an average number of 31 water molecules, and this number can change and diminish substantially upon increase in temperature and decrease in density. To explain the unexpected inverse fractionation observed for benzene, one could hypothesize a similar effect on benzene hydration due to the deuterium substitutions. Processes connected to the hydrophobic effect of aromatic molecules in solution and associated changes of density and orientation of water molecules surrounding the solute,<sup>51</sup> as well as the structure and patterns of solute hydration,<sup>52</sup> appear to be more important than the effects due to mass and molar volume differences. This hypothesis will require future substantiation that might be provided by molecular dynamic simulations of aqueous diffusion of deuterated and nondeuterated species. Such simulations have been performed, for instance, for charged species<sup>22,53,54</sup> and have contributed to elucidate the importance of the hydration shell and of the interaction between solute and water molecules for the observed isotope fractionation of different ions in aqueous solution. The different interaction of deuterated and nondeuterated benzene and toluene molecules with water molecules appears to be an important factor to explain the experimental observations of this study. This can be deduced also by comparing the results of this study with earlier experiments. For instance, benzene self-diffusion<sup>55</sup> and tracer diffusion in chlorobenzene<sup>56</sup> showed only very minor or no isotope effects during diffusion of  $C_6H_6$  and  $C_6D_6$  in these organic liquids, which contrasts with the rather large fractionation consistently observed in the different aqueous systems considered in our study.

We think that the normal and inverse diffusive fractionation effects observed for deuterated and nondeuterated BTEX will stimulate further research to develop a mechanistic understanding of transport and transformation of these contaminants and their labeled mixtures in different environments. Despite fractionation of D/H, substituted aromatic compounds due to biodegradation can be large and may be dominant for many transformation pathways;<sup>57</sup> our study shows that diffusive fractionation is also significant and should be considered when these chemicals are used as tracers and diagnostic tools in environmental systems in which transport and transformation processes are typically coupled. On a broader perspective, the results of this study, showing different diffusive isotope fractionation of very similar compounds, also have implications for the fast developing field of compound-specific stable isotope analysis (CSIA) techniques,<sup>58,59</sup> which will greatly benefit from data for different environmental contaminants. The path toward developing a comprehensive database of organic pollutant diffusive isotope fractionation and a sound theory capturing the key mechanisms causing diffusive fractionation is inherently complex but rich of intriguing future challenges.

## ■ ASSOCIATED CONTENT

### § Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.estlett.7b00159.

Descriptions of the flow-through system and modeling approach used to quantitatively interpret the results in the different experimental setups. (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*Phone: +45 45251566. E-mail: masro@env.dtu.dk.

### ORCID

Massimo Rolle: 0000-0001-8833-8951

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

The authors acknowledge the support of the Deutsche Forschungsgemeinschaft (Grant RO 4169/2-1) and the help of Chrysanthi-Elisabeth Nika, Hanne Bøggild, and Mikael Emil Olsson for assistance in the experimental work.

## ■ REFERENCES

- (1) Wiedemeier, T. H.; Rifai, H. S.; Newell, C. H.; Wilson, T. H. *Natural Attenuation of Fuels and Chlorinated Solvents in the Subsurface*; John Wiley: New York, 1999.
- (2) Thierrin, J.; Davis, G. B.; Barber, C. A Groundwater Tracer Test with Deuterated Compounds for Monitoring In-Situ Biodegradation and Retardation of Aromatic Hydrocarbons. *Groundwater* **1995**, *33*, 469–475.
- (3) Reusser, D. E.; Istok, J. D.; Beller, H. R.; Field, J. A. In situ transformation of deuterated toluene and xylene to benzylsuccinic acid analogues in BTEX-contaminated aquifers. *Environ. Sci. Technol.* **2002**, *36*, 4127–4134.
- (4) Fischer, A.; Bauer, J.; Meckenstock, R. U.; Stichler, W.; Griebler, C.; Maloszewski, P.; Kastner, M.; Richnow, H. H. A multitracer test proving the reliability of Rayleigh equation-based approach for assessing biodegradation in a BTEX contaminated aquifer. *Environ. Sci. Technol.* **2006**, *40*, 4245–4252.
- (5) Gieg, L. M.; Alumbaugh, R. E.; Field, J.; Jones, J.; Istok, J. D.; Suflita, J. M. Assessing in situ rates of anaerobic hydrocarbon bioremediation. *Microb. Biotechnol.* **2009**, *2*, 222–233.
- (6) Cozzarelli, I. M.; Bekins, B. A.; Eganhouse, R. P.; Warren, E.; Essaid, H. I. In situ measurements of volatile aromatic hydrocarbon biodegradation rates in groundwater. *J. Contam. Hydrol.* **2010**, *111*, 48–64.
- (7) Parker, B. L.; Chapman, S. W.; Guilbeault, M. A. Plume persistence caused by back diffusion from thin clay layers in a sand aquifer following TCE source-zone hydraulic isolation. *J. Contam. Hydrol.* **2008**, *102*, 86–104.
- (8) Carrera, J.; Sanchez-Vila, X.; Benet, I.; Medina, A.; Galarza, G.; Guimera, J. On matrix diffusion: formulations, solution methods and qualitative effects. *Hydrogeol. J.* **1998**, *6*, 178–90.
- (9) Rolle, M.; Chiogna, G.; Hochstetler, D. L.; Kitanidis, P. K. On the importance of diffusion and compound-specific mixing for groundwater transport: An investigation from pore to field scale. *J. Contam. Hydrol.* **2013**, *153*, 51–68.
- (10) Rolle, M.; Kitanidis, P. K. Effects of compound-specific dilution on transient transport and solute breakthrough: A pore-scale analysis. *Adv. Water Resour.* **2014**, *71*, 186–199.
- (11) LaBolle, E. M.; Fogg, G. E.; Eweis, J. B.; Gravner, J.; Leaist, D. G. Isotopic fractionation by diffusion in groundwater. *Water Resour. Res.* **2008**, *44*, W07405.
- (12) Rolle, M.; Chiogna, G.; Bauer, R.; Griebler, C.; Grathwohl, P. Isotopic Fractionation by Transverse Dispersion: Flow-through

Microcosms and Reactive Transport Modeling Study. *Environ. Sci. Technol.* **2010**, *44*, 6167–6173.

(13) Eckert, D.; Rolle, M.; Cirpka, O. A. Numerical simulation of isotope fractionation in steady-state bioreactive transport controlled by transverse mixing. *J. Contam. Hydrol.* **2012**, *140-141*, 95–106.

(14) Van Breukelen, B. M.; Rolle, M. Transverse Hydrodynamic Dispersion Effects on Isotope Signals in Groundwater Chlorinated Solvents' Plumes. *Environ. Sci. Technol.* **2012**, *46*, 7700–7708.

(15) Jin, B.; Rolle, M.; Li, T.; Haderlein, S. B. Diffusive fractionation of BTEX and chlorinated ethenes in aqueous solution: quantification of spatial isotope gradients. *Environ. Sci. Technol.* **2014**, *48*, 6141–6150.

(16) Wanner, P.; Hunkeler, D. Carbon and chlorine isotopologue fractionation of chlorinated hydrocarbons during diffusion in water and low permeability sediments. *Geochim. Cosmochim. Acta* **2015**, *157*, 198–212.

(17) Jähne, B.; Heinz, G.; Dietrich, W. Measurement of the diffusion coefficients of sparingly soluble gases in water. *J. Geophys. Res.* **1987**, *92*, 10767–10776.

(18) Eggenkamp, H. G. M.; Middelburg, J. J.; Kreulen, R. Preferential diffusion of  $^{35}\text{Cl}$  relative to  $^{37}\text{Cl}$  in sediments of Kau Bay, Halmahera, Indonesia. *Chem. Geol.* **1994**, *116*, 317–325.

(19) Prinzhofer, A.; Pernaton, E. Isotopically light methane in natural gas: bacterial imprint or diffusive fractionation? *Chem. Geol.* **1997**, *142*, 193–200.

(20) Schloemer, S.; Krooss, B. M. Molecular transport of methane, ethane and nitrogen and the influence of diffusion on the chemical and isotopic composition of natural gas accumulations. *Geofluids* **2004**, *4*, 81–108.

(21) Richter, F. M.; Mendybaev, R. A.; Christensen, J. N.; Hutcheon, I. D.; Williams, R. W.; Sturchio, N. C.; Beloso, A. D. Kinetic isotopic fractionation during diffusion of ionic species in water. *Geochim. Cosmochim. Acta* **2006**, *70*, 277–289.

(22) Bourq, I. C.; Sposito, G. Molecular dynamics simulations of kinetic isotope fractionation during the diffusion of ionic species in liquid water. *Geochim. Cosmochim. Acta* **2007**, *71*, 5583–5589.

(23) Donahue, M. A.; Werne, J. P.; Meile, C.; Lyons, T. W. Modeling sulfur isotope fractionation and differential diffusion during sulfate reduction in sediments of the Cariaco Basin. *Geochim. Cosmochim. Acta* **2008**, *72*, 2287–2297.

(24) Bourq, I. C.; Sposito, G. Isotopic fractionation of noble gases by diffusion in liquid water: Molecular dynamics simulations and hydrologic applications. *Geochim. Cosmochim. Acta* **2008**, *72*, 2237–2247.

(25) Eggenkamp, H. G. M.; Coleman, M. L. The effect of aqueous diffusion on the fractionation of chlorine and bromine stable isotopes. *Geochim. Cosmochim. Acta* **2009**, *73*, 3539–3548.

(26) Bourq, I. C.; Richter, F. M.; Christensen, J. N.; Sposito, G. Isotopic mass dependence of metal cation diffusion coefficients in liquid water. *Geochim. Cosmochim. Acta* **2010**, *74*, 2249–2256.

(27) Beekman, H. E.; Eggenkamp, H. G. M.; Appelo, C. A. J. An integrated modelling approach to reconstruct complex solute transport mechanisms - Cl and delta Cl-37 in pore water of sediments from a former brackish lagoon in The Netherlands. *Appl. Geochem.* **2011**, *26*, 257–268.

(28) Wortmann, U. G.; Chernyavsky, B. M. The significance of isotope specific diffusion coefficients for reaction-transport models of sulfate reduction in marine sediments. *Geochim. Cosmochim. Acta* **2011**, *75*, 3046–3056.

(29) Tempest, K.; Emerson, S. Kinetic isotope fractionation of argon and neon in during air–water transfer. *Mar. Chem.* **2013**, *153*, 39–47.

(30) Richter, F. M.; Liang, Y.; Davis, A. M. Isotope fractionation by diffusion in molten oxides. *Geochim. Cosmochim. Acta* **1999**, *63*, 2853–2861.

(31) Richter, F. M.; Davis, A. M.; DePaolo, D. J.; Watson, E. B. Isotope fractionation by chemical diffusion between molten basalt and rhyolite. *Geochim. Cosmochim. Acta* **2003**, *67*, 3905–3923.

(32) Watkins, M. W.; DePaolo, D. J.; Ryerson, F. J.; Peterson, B. T. Influence of liquid structure on diffusive isotope separation in molten

silicates and aqueous solutions. *Geochim. Cosmochim. Acta* **2011**, *75*, 3103–3118.

(33) Watkins, M. W.; Liang, Y.; Richter, F. M.; Ryerson, F. J.; DePaolo, D. J. Diffusion of multi-isotopic chemical species in molten silicates. *Geochim. Cosmochim. Acta* **2014**, *139*, 313–326.

(34) Mills, R. Search for isotope effects in the self-diffusion of benzene and cyclohexane at 25°. *J. Phys. Chem.* **1975**, *79*, 852–853.

(35) Mills, R.; Harris, K. R. The effect of isotopic substitution on diffusion in liquids. *Chem. Soc. Rev.* **1976**, *5*, 215–231.

(36) Freer, R.; Sherwood, J. N. Diffusion in organic liquids. *J. Chem. Soc., Faraday Trans. 1* **1980**, *76*, 1030–1037.

(37) Haberer, C. M.; Rolle, M.; Cirpka, O. A.; Grathwohl, P. Oxygen transfer in a fluctuating capillary fringe. *Vadose Zone J.* **2012**, *11*, 0.

(38) Cussler, E. L. *Diffusion–Mass Transfer in Fluid Systems*, 3rd ed.; Cambridge University Press, 2009.

(39) Cirpka, O. A.; Rolle, M.; Chiogna, G.; de Barros, F. P. J.; Nowak, W. Stochastic evaluation of mixing-controlled steady-state plume lengths in two-dimensional heterogeneous domains. *J. Contam. Hydrol.* **2012**, *138-139*, 22–39.

(40) Muniruzzaman, M.; Haberer, C. M.; Grathwohl, P.; Rolle, M. Multicomponent ionic dispersion during transport of electrolytes in heterogeneous porous media: Experiments and model-based interpretation. *Geochim. Cosmochim. Acta* **2014**, *141*, 656–669.

(41) Bhattacharyya, S.; Bagchi, B. Power law mass dependence of diffusion: A mode coupling theory analysis. *Phys. Rev. E: Stat. Phys., Plasmas, Fluids, Relat. Interdiscip. Top.* **2000**, *61*, 3850–3856.

(42) Ye, Y.; Chiogna, G.; Cirpka, O. A.; Grathwohl, P.; Rolle, M. Experimental investigation of compound-specific dilution of solute plumes in saturated porous media: 2-D vs. 3-D flow-through systems. *J. Contam. Hydrol.* **2015**, *172*, 33–47.

(43) Chiogna, G.; Eberhardt, C.; Grathwohl, P.; Cirpka, O. A.; Rolle, M. Evidence of compound-dependent hydrodynamic and mechanical transverse dispersion by multitracer laboratory experiments. *Environ. Sci. Technol.* **2010**, *44*, 688–693.

(44) Rolle, M.; Muniruzzaman, M.; Haberer, C. M.; Grathwohl, P. Coulombic effects in advection-dominated transport of electrolytes in porous media: Multicomponent ionic dispersion. *Geochim. Cosmochim. Acta* **2013**, *120*, 195–205.

(45) Muniruzzaman, M.; Rolle, M. Impact of multicomponent ionic transport on pH fronts propagation in saturated porous media. *Water Resour. Res.* **2015**, *51*, 6739–6755.

(46) Rolle, M.; Hochstetler, D. L.; Chiogna, G.; Kitanidis, P. K.; Grathwohl, P. Experimental investigation and pore-scale modeling interpretation of compound-specific transverse dispersion in porous media. *Transp. Porous Media* **2012**, *93*, 347–362.

(47) Hochstetler, D. L.; Rolle, M.; Chiogna, G.; Haberer, C. M.; Grathwohl, P.; Kitanidis, P. K. Effects of compound-specific transverse mixing on steady-state reactive plumes: Insights from pore-scale simulations and Darcy-scale experiments. *Adv. Water Resour.* **2013**, *54*, 1–10.

(48) Bartell, L. S.; Roskos, R. R. Isotope effects on molar volume and surface tension: simple theoretical model and experimental data for hydrocarbons. *J. Chem. Phys.* **1966**, *44*, 457–463.

(49) Dunitz, J.; Ibberson, R. M. Is deuterium always smaller than protium? *Angew. Chem.* **2008**, *120*, 4276–4278.

(50) Choudhary, A.; Chandra, A. Spatial and orientational structure of the hydration shell of benzene in sub- and supercritical water. *J. Phys. Chem. B* **2015**, *119*, 8600–8612.

(51) Raschke, T.; Levitt, M. Nonpolar solutes enhance water structure within hydration shells while reducing interactions between them. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 6777–6782.

(52) Schravendijk, P.; van der Vegt, N. F. A. From hydrophobic to hydrophilic solvation: An application to hydration of benzene. *J. Chem. Theory Comput.* **2005**, *1*, 643–652.

(53) Moller, K. B.; Rey, R.; Masia, M.; Hynes, J. T. On the coupling between molecular diffusion and solvation shell exchange. *J. Chem. Phys.* **2005**, *122*, 114508.

(54) Hofmann, A. E.; Bourg, I. C.; DePaolo, D. J. Ion desolvation as a mechanism for kinetic isotope fractionation in aqueous systems. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 18689–18694.

(55) Mills, R. Diffusion relationships in the binary system benzene-perdeuteriobenzene at 25 C. *J. Phys. Chem.* **1976**, *80*, 888–890.

(56) Shankland, I. R.; Dunlop, P. J.; Barr, L. W. Isotope effect in liquids as shown by tracer diffusion at 25 C of several labeled benzenes in benzene, chlorobenzene, n-octane, and cyclohexane, and the relationship to the isotope effect in solids. *Phys. Rev. B* **1975**, *12*, 2249–2252.

(57) Morasch, B.; Richnow, H.-H.; Schink, B.; Meckenstock, R. U. Stable hydrogen and carbon isotope fractionation during microbial toluene degradation: mechanistic and environmental aspects. *Appl. Environ. Microbiol.* **2001**, *67*, 4842–4849.

(58) Thullner, M.; Centler, F.; Richnow, H. H.; Fischer, A. Quantification of organic pollutant degradation in contaminated aquifers using compound specific stable isotope analysis - Review of recent developments. *Org. Geochem.* **2012**, *42*, 1440–1460.

(59) Hatzinger, P. B.; Bohlke, J.; Sturchio, N. C. Application of stable isotope ratio analysis for biodegradation monitoring in groundwater. *Curr. Opin. Biotechnol.* **2013**, *24*, 542–549.